

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

INTAKE[®] COMBI IN-FURROW AND FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250g/L FLUTRIAFOL

GROUP	3	FUNGICIDE
-------	----------	-----------

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the directions for use table.

CONTENTS: 10, 110, 200 & 1000 LITRES

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

DIRECTIONS FOR USE : ALL STATES

Restrictions: DO NOT use open loading and mixing for aerial application.

A. INTAKE COMBI APPLIED IN-FURROW AS A FERTILISER TREATMENT

Crop	Disease	Rate per hectare	WHP	Critical Comments												
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400mL	4 weeks Grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table data-bbox="1196 523 2047 730"> <thead> <tr> <th data-bbox="1196 523 1563 582"><u>Fertiliser rate per hectare (kg)</u></th> <th data-bbox="1576 523 2047 582"><u>Volume of product per 100kg of fertiliser (mL)</u></th> </tr> </thead> <tbody> <tr> <td data-bbox="1279 587 1368 614">60</td> <td data-bbox="1778 587 1839 614">667</td> </tr> <tr> <td data-bbox="1279 619 1368 646">70</td> <td data-bbox="1778 619 1839 646">571</td> </tr> <tr> <td data-bbox="1279 651 1368 678">80</td> <td data-bbox="1778 651 1839 678">500</td> </tr> <tr> <td data-bbox="1279 683 1368 710">100</td> <td data-bbox="1778 683 1839 710">400</td> </tr> <tr> <td data-bbox="1279 715 1368 742">120</td> <td data-bbox="1778 715 1839 742">333</td> </tr> </tbody> </table>	<u>Fertiliser rate per hectare (kg)</u>	<u>Volume of product per 100kg of fertiliser (mL)</u>	60	667	70	571	80	500	100	400	120	333
<u>Fertiliser rate per hectare (kg)</u>	<u>Volume of product per 100kg of fertiliser (mL)</u>															
60	667															
70	571															
80	500															
100	400															
120	333															

Crop	Disease	Rate per hectare	WHP	Critical Comments																																							
Wheat	Stripe rust (<i>Puccinia striiformis</i> f.sp. <i>tritici</i>)	200 to 800mL	4 weeks Grazing	<p>For stripe rust control in wheat use 200 to 400mL/ha where up to 100 days protection is required and where heavy stripe rust pressure is expected use 400mL/ha only. Use 600 to 800mL/ha when 120 to 130 days protection is required and where heavy stripe rust pressure is expected, use 800mL/ha only.</p> <p>For control of powdery mildew and scald of barley, use 200 to 400mL/ha where up to 80 days protection is required and where heavy disease pressure is expected use 400mL/ha. Use 600mL/ha when 100 days protection is required and 800mL/ha for up to 120 days protection from powdery mildew and scald.</p> <p>For suppression of net form net blotch for up to 110 days after sowing use 600 to 800mL/ha.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2">Fertiliser rate per hectare (kg)</th> <th colspan="4">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <th>200mL rate</th> <th>400mL rate</th> <th>600mL rate</th> <th>800mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>400</td> <td>800*</td> <td>1200*</td> <td>1600*</td> </tr> <tr> <td>60</td> <td>333</td> <td>667*</td> <td>1000*</td> <td>1332*</td> </tr> <tr> <td>70</td> <td>286</td> <td>571</td> <td>858*</td> <td>1114*</td> </tr> <tr> <td>80</td> <td>250[†]</td> <td>500</td> <td>750*</td> <td>1000*</td> </tr> <tr> <td>100</td> <td>200[†]</td> <td>400</td> <td>600</td> <td>800*</td> </tr> <tr> <td>120</td> <td>167[†]</td> <td>333</td> <td>500</td> <td>668*</td> </tr> </tbody> </table> <p>* Application volumes over 600mL per 100kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. [†] Where application volume is low, product may be diluted with water to improve coverage.</p>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)				200mL rate	400mL rate	600mL rate	800mL rate	50	400	800*	1200*	1600*	60	333	667*	1000*	1332*	70	286	571	858*	1114*	80	250 [†]	500	750*	1000*	100	200 [†]	400	600	800*	120	167 [†]	333	500	668*
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																																									
200mL rate		400mL rate	600mL rate	800mL rate																																							
50	400	800*	1200*	1600*																																							
60	333	667*	1000*	1332*																																							
70	286	571	858*	1114*																																							
80	250 [†]	500	750*	1000*																																							
100	200 [†]	400	600	800*																																							
120	167 [†]	333	500	668*																																							
	Septoria tritici blotch (<i>Mycosphaerella graminicola</i>) Take-all (<i>Geaumannomyces graminis</i>)	400mL																																									
Barley	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>hordei</i>) Scald (<i>Rhynchosporium secalis</i>)	200 to 800mL																																									
	Net Form Net Blotch (<i>Pyrenophora teres</i> f.sp. <i>teres</i>) (Suppression only)	600 to 800mL																																									

B. INTAKE COMBI APPLIED AS A FOLIAR SPRAY

Crop	Disease	Rate per hectare	WHP	Critical Comments
Wheat	Stripe rust (<i>Puccinia striiformis</i> f.sp. <i>tritici</i>)	250mL - 500mL Ground Application Apply in 100 - 150L water plus 200mL 'BS 1000' per 100L of spray solution.	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.
	Leaf rust (<i>P. recondita</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>) Septoria tritici blotch (<i>Mycosphaerella graminicola</i>)	Aerial Application Apply in a minimum of 15L water plus 200mL 'BS 1000' per 100L of spray solution.		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.
Barley	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>hordei</i>)	Apply in a minimum of 15L water plus 200mL 'BS 1000' per 100L of spray solution.	10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate, may be required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

A. APPLIED IN-FURROW AS A FERTILISER TREATMENT :

CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. APPLIED AS A FOLIAR SPRAY:

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing : Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.

Foliar Application

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

Compatibility

INTAKE COMBI is compatible with the liquid fertilizers – Flexi N[®], Flexi NK[®], Flexi NS[®] and UAN. Pre mixing of INTAKE COMBI with water (0.5:1 to 1:1) is recommended. Agitation of the liquid fertilizer prior to and during addition of INTAKE COMBI is required. Maintain agitation of the mix during application.

Re-entry or Re-handling

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	----------	-----------

INTAKE COMBI IN-FURROW AND FOLIAR FUNGICIDE is a member of the DMI group of fungicides. For fungicide resistance management INTAKE COMBI IN-FURROW AND FOLIAR FUNGICIDE is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to INTAKE COMBI IN-FURROW AND FOLIAR FUNGICIDE and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by INTAKE COMBI IN-FURROW AND FOLIAR FUNGICIDE or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of INTAKE COMBI IN-FURROW AND FOLIAR FUNGICIDE to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (10 & 200L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

STORAGE AND DISPOSAL (110 & 1000L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Crop Care Australasia Pty Ltd should be advised immediately.

This container is reusable and remains the property of Crop Care Australasia Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Crop Care Australasia Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing the spray wear elbow length chemical resistant gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier or from the Crop Care website: www.cropcare.com.au

Condition of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS – AUSTRALIA WIDE
---	--

'Intake', 'Flexi-N', 'Flexi-NK' and 'Flexi NS' are Registered Trademarks.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie Qld 4172
Tel : (07) 3909 2000

Insert barcodes,
label code and
drumMUSTER
logo as required

APVMA Approval No. : 61702/50423

BN
DOM