

POISON


KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Paramectin[®]

POUR-ON FOR CATTLE

ACTIVE CONSTITUENTS: 5 g/L ABAMECTIN
SOLVENT: 950 g /L DIETHYLENE GLYCOL
MONOBUTYL ETHER

For the treatment and control of abamectin sensitive
internal and external parasites of cattle.



1L

Jurox

406507

Front Label

PEEL
HERE

Paramectin™ Pour-On for Cattle

READ THE ENCLOSED INFORMATION BEFORE USING THIS PRODUCT

SAFETY DIRECTIONS: Poisonous if absorbed by skin contact or swallowed. Will damage eyes. May irritate the nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When using the product wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and goggles. If clothing becomes contaminated with the product, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day@use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre, phone 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

STORE below 30°C (room temperature). Store in original container.

APVMA 50341/1L/1004



Jurox Pty Ltd
85 Gardiners Road
Rutherford NSW 2320
Infoline 1800 023 312

Ⓟ

EXP.

406512



9 338094 000158 >

PARAMECTIN® Pour-On for Cattle

INDICATIONS:

Paramectin Pour-On for Cattle contains abamectin, a member of the avermectin family of compounds. It provides a broad spectrum of efficacy against sensitive strains of the following adult and immature internal and external parasites of cattle. Resistance may develop to any drench.

Adult and immature gastrointestinal roundworms

- *Haemonchus placei* - Barber's pole worm
- *Cooperia oncophora*, *C. punctata* - Small intestinal worm
- *Ostertagia ostertagi* - Small brown stomach worm (including inhibited L4 stage). Controls infection acquired up to 14 days after treatment.
- *Nematodirus spathiger* - Thin-necked intestinal worm
- *Oesophagostomum radiatum* - Nodule worm
- *Oesophagostomum venulosum* - Large bowel worm
- *Trichostrongylus axei* - Stomach hair worm
- *Trichuris* spp - Whip worm
- *Trichostrongylus colubriformis* - Intestinal hair worm
- *Stongyloides papillosus* - Intestinal threadworm

Lungworms

- *Dictyocaulus viviparus* - Large lungworm.
Controls infection acquired up to 28 days after treatment.

Sucking Lice and Biting Lice

- *Linognathus vituli* - Long-nosed sucking louse
- *Haematopinus eurysternus* - Short-nosed sucking louse
- *Bovicola bovis* - Biting louse
- *Solenopotes capillatus* - Little blue sucking louse

Cattle Tick *Boophilus microplus*

Buffalo Fly *Haematobia irritans exigua*. Controls buffalo fly and protects cattle from reinfestation for up to 14 days after treatment.

PARAMECTIN® Pour-On is not recommended in strategic dipping programmes.

DIRECTIONS FOR USE:

DO NOT treat calves under 50 kg.

The minimum retreatment interval permitted is 35 days.

Dose the mob according to the heaviest animal by liveweight in the group (bulls, cows, steers, calves etc). Do not underdose. A representative sample of animals should be weighed before treatment either with scales or a weighband. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Effect on Dung Beetles

When used according to the label directions, Paramectin Pour-On for Cattle is not likely to have any unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations of the following species: *Onthophagus australis*, *O. granulatus*, *O. dandalu*, *O. chepara*, *Onitis alexis*, *Euoniticellus intermedius*, *E. africanus* and *Apodius* spp. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.

DOSE: 1 mL per 10 kg bodyweight (500 µg/kg bodyweight).

Liveweight	Dose	1 L treats
50 - 75 kg	7.5 mL	133
76 - 100 kg	10 mL	100
101 - 150 kg	15 mL	66
151 - 200 kg	20 mL	50
201 - 250 kg	25 mL	40
251 - 300 kg	30 mL	33
301 - 350 kg	35 mL	28
351 - 400 kg	40 mL	25
401 - 450 kg	45 mL	22
451 - 500 kg	50 mL	20
501 - 550 kg	55 mL	18
551 - 600 kg	60 mL	16
601 - 650 kg	65 mL	15

Cattle in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 35 days before slaughter for human consumption.

MILK: Nil

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT SLAUGHTER for export within 42 days of treatment.

SAFETY DIRECTIONS: Poisonous if absorbed by skin contact or swallowed. Will damage eyes. May irritate the nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When using the product wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and goggles. If clothing becomes contaminated with the product, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre, phone 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT: Abamectin is extremely toxic to aquatic species. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.

PARAMECTIN® Pour-On for Cattle

INDICATIONS:

Paramectin Pour-On for Cattle contains abamectin, a member of the avermectin family of compounds. It provides a broad spectrum of efficacy against sensitive strains of the following adult and immature internal and external parasites of cattle. Resistance may develop to any drench.

Adult and immature gastrointestinal roundworms

¥*Haemonchus placei* - Barber's pole worm

¥*Cooperia oncophora*, *C. punctata* - Small intestinal worm

¥*Ostertagia ostertagi* - Small brown stomach worm (including inhibited L4 stage). Controls infection acquired up to 14 days after treatment.

¥*Nematodirus spathiger* - Thin-necked intestinal worm

¥*Oesophagostomum radiatum* - Nodule worm

¥*Oesophagostomum venulosum* - Large bowel worm

¥*Trichostrongylus axei* - Stomach hair worm

¥*Trichuris* spp - Whip worm

¥*Trichostrongylus colubriformis* - Intestinal hair worm

¥*Stongyloides papillosus* - Intestinal threadworm

Lungworms

¥*Dictyocaulus viviparus* - Large lungworm.

Controls infection acquired up to 28 days after treatment.

Sucking Lice and Biting Lice

¥*Linognathus vituli* - Long-nosed sucking louse

¥*Haematopinus eurysternus* - Short-nosed sucking louse

¥*Bovicola bovis* - Biting louse

¥*Solenopotes capillatus* - Little blue sucking louse

Cattle Tick *Boophilus microplus*

Buffalo Fly *Haematobia irritans exigua*. Controls buffalo fly and protects cattle from reinfestation for up to 14 days after treatment.

PARAMECTIN® Pour-On is not recommended in strategic dipping programmes.

DIRECTIONS FOR USE:

DO NOT treat calves under 50 kg.

The minimum retreatment interval permitted is 35 days.

Dose the mob according to the heaviest animal by liveweight in the group (bulls, cows, steers, calves etc). Do not underdose. A representative sample of animals should be weighed before treatment either with scales or a weighband. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Effect on Dung Beetles

When used according to the label directions, Paramectin Pour-On for Cattle is not likely to have any unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations of the following species: *Onthophagus australis*, *O. granulatus*, *O. dandalu*, *O. chepara*, *Onitis alexis*, *Euoniticellus intermedius*, *E. africanus* and *Apodius* spp. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.

DOSE: 1 mL per 10 kg bodyweight (500 µg/kg bodyweight).

Liveweight	Dose	1 L treats
50 - 75 kg	7.5 mL	133
76 - 100 kg	10 mL	100
101 - 150 kg	15 mL	66
151 - 200 kg	20 mL	50
201 - 250 kg	25 mL	40
251 - 300 kg	30 mL	33
301 - 350 kg	35 mL	28
351 - 400 kg	40 mL	25
401 - 450 kg	45 mL	22
451 - 500 kg	50 mL	20
501 - 550 kg	55 mL	18
551 - 600 kg	60 mL	16
601 - 650 kg	65 mL	15

Cattle in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 35 days before slaughter for human consumption.

MILK: Nil

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT SLAUGHTER for export within 42 days of treatment.

SAFETY DIRECTIONS: Poisonous if absorbed by skin contact or swallowed. Will damage eyes. May irritate the nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When using the product wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and goggles. If clothing becomes contaminated with the product, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day of use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre, phone 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE

ENVIRONMENT: Abamectin is extremely toxic to aquatic species. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.