



## DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
 CAN KILL IF SWALLOWED  
 DO NOT PUT IN DRINK BOTTLES  
 KEEP LOCKED UP

Nufarm  
**Nuquat<sup>®</sup> 250**  
 Non-Residual Knockdown Herbicide

**ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE**

GROUP **L** HERBICIDE

For the control of a wide range of grasses and broadleaf weeds as per the directions for use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

© Nuquat is a registered trademark of Nufarm Australia Limited

[nufarm.com.au](http://nufarm.com.au)

Nufarm Australia Limited  
 ACN 004 377 780  
 103-105 Pipe Road  
 Laverton North Victoria 3026  
 Tel: (03) 9282 1000  
 Fax: (03) 9282 1001

Contents: 20 Litres  
 Contents: 110 Litres  
 Contents 1000Litres

APVMA Approval No.: 53221/54620

**FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.  
 THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.**

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should NOT be burnt.

### **Envirodrum 110L Mini Bulk Returnable Container**

Store in the original sealed Envirodrum in a cool, well ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited. (1000L)

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks eyes. Protect eyes while using. Avoid contact with eyes, skin and clothing. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use wash contaminated clothing, gloves and face shield or goggles.

**SPRAY APPLICATION**

- DO NOT work in spray mist.
- DO NOT continue to use if skin irritation or nose bleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
- When using misting machines for weed control in banana plantations, cut back to run at half throttle, thus preventing the production of fine droplets, the inhalation of which may be dangerous.
- When there is a risk of exposure to spray mist or when using misting machines in banana plantations wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended, but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
- Avoid contacting vegetation wet with spray, but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

**FIRST AID**

If poisoning occurs, get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**NOTE TO PHYSICIANS**

For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning; A Guide For Doctors".

**MATERIAL SAFETY DATA SHEET**


For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from you supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

BN / DOM / Barcode / /

<p><b>BIPYRIDILIUM PESTICIDES, LIQUID, TOXIC, (contains paraquat)</b></p>	
<p><b>UN No: 3016</b></p>	
<p><b>PACKING GROUP: III</b></p>	
<p><b>HAZCHEM: 2 X</b></p>	
<p><b>IN AN EMERGENCY DIAL: 000 POLICE OR FIRE BRIGADE</b></p>	<p><b>IN AN EMERGENCY, SPECIALIST ADVICE NUFARM AUSTRALIA LIMITED 1800 033 498</b></p>



# DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
CAN KILL IF SWALLOWED  
DO NOT PUT IN DRINK BOTTLES  
KEEP LOCKED UP

## Nufarm Nuquat<sup>®</sup> 250

Non-Residual, Knockdown Herbicide

**ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE**

GROUP **L** HERBICIDE

For the control of a wide range of grasses and broadleaf weeds as per the directions for use.

READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

© Nuquat is a registered trademark of Nufarm Australia Limited

nufarm.com.au

APVMA Approval No.: 53221/54620

### DIRECTIONS FOR USE

#### RESTRAINTS

DO NOT add wetter unless spraying at high volume. Where Nuquat 250 is mixed with water at less than 400mL/100L of water, add 70mL Activator<sup>®</sup> or 60mL Chemwet 1000 per 100L of spray.

DO NOT spray plants which are waterlogged, under stress of any kind or covered with soil or dust.

DO NOT spray plants covered with heavy dew, but rain following spraying will not affect results.

DO NOT sow or cultivate for 1 hour after spraying but operations should commence within 7 days.

**For ground application only** - DO NOT use through aircraft, misting machines (except in banana plantations) or hand-held ultra low volume controlled droplet applicators (CDA units).

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS	
Rice	Annual grass and broadleaf weed control.  Early Autumn sowing.	All States	1.2-1.6L/ha <sup>†</sup>	Where cultivation follows spraying, it may commence one hour after spraying but should be completed within 7 days. Where heavy weed growth is present at spraying a better seed bed will result if cultivation is delayed 3-5 days.	
	Winter, Spring and early Summer sowing		1.6-2.4L/ha		
	Volunteer canola including Roundup Ready <sup>®</sup> varieties (up to 4 leaf) (up to 6 leaf)		1.8L/ha 2.4L/ha		
	Wild oats at 2-5 leaf stage in Autumn/Winter.	Qld, Vic, SA, Tas, NT, WA only	600-800mL/ha	Pre-sowing. Post-sowing, pre-crop emergence.	
		NSW, ACT only	600mL/ha		
Annual grass and broadleaf weed control.	Qld, NSW, NT only	1.6L/ha 800mL/ha			
Wild Oat control in Spring Fallows	Wild oats at 2-5 leaf stage	Qld, NSW, NT, ACT only	1.2-2L/ha		Use higher rate for Summer growth. Avoid spraying under hot, dry conditions. Best results will be obtained when spraying is carried out in the late evening.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Kikuyu/ Paspalum Pasture	To suppress growth to oversow Winter seed.	Qld, NSW, ACT only	1.6 or 2.4L/ha	Use the high rate for February spraying and the low rate in March
Selective Weed Control Autumn/ early Winter - annual clovers - perennial clover  Late Winter/early Spring - annual clovers - perennial clovers - cocksfoot - perennial ryegrass - phalaris - Demeter fescue only	Annual grass and some broadleaf weed control except Paterson's curse, Sorrel, Dock, Shepherd's purse and some thistles	All States	600mL-1.2L/ha  1.2-1.6L/ha <sup>‡</sup>	Use the higher rates for dense weed stands.
	For control of these weeds alternative methods such as the spray-graze technique with Amicide® 625 or MCPA should be considered.	Qld, NSW, Vic, Tas, NT, SA, ACT only	1.6-2.4L/ha <sup>‡</sup>	Use the higher rate in Winter-early Spring when barley grass is present. <b>All applications:</b> Graze pastures continuously after the seasonal break to a height of 2-4cm. Remove stock 2-3 days before spraying to allow weeds to freshen up. DO NOT apply until clover has reached the 6 leaf stage. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following Spring rainfall and growth. DO NOT spray clovers which are affected by insect attack, disease or moisture stress and DO NOT use on clover pastures growing in water repellent sands or other situations subject to moisture stress at or immediately following treatment otherwise poor recovery of the clover may result. Use the lower rate for cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter Fescue. The perennial grasses must be at least 12 months old at spraying. <b>DO NOT APPLY TO MEDICS</b>
	Yorkshire fog grass		1.2L/ha	Apply in early Spring to reduce Yorkshire fog grass component and increase the clover and desirable grass component. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following Spring rainfall and growth. In lower rainfall areas application in mid to late Winter may be almost as effective but allow better pasture recovery. If pasture has been grazed, allow sufficient time for pasture and fog grass recovery before spraying. Apply in spray volumes of 100-250L/ha, the latter for dense or tall, ungrazed pastures. Add Activator® at 140mL/100L or Chemwet 1000 at 120mL/100L.
Lucerne Autumn/ early Winter  Late Winter/ early Spring	Annual grass and some broadleaf weeds	Qld, Vic, SA, WA, Tas, NT only	1.2-1.6L/ha <sup>‡</sup>	Use the higher rates for dense weed stands. DO NOT spray Lucerne stands under 12 months old. For residual weed control or if Paterson's curse, Shepherd's purse and some other broadleaf weeds are present add Nufarm Diuron 900DF Herbicide at 1.1kg or 1.9kg or Nufarm Flowable Diuron at 2L or 3.5L/ha. If mintweed is present use Nufarm Nu-Trazine 900DF Herbicide at 600g/ha or Nufarm Flowable Nu-trazine at 1.1L/ha.  <b>WARNING</b> - In certain areas, an uncommon species of barley grass ( <i>H. glaucum</i> - common barley grass is <i>H. leporinum</i> ) resistant to paraquat based products has become established. It may regrow after an initial scorch by Nuquat 250. Where the problem is suspected use a suitable post-emergent grass herbicide (such as Nufarm Asset®) at recommended label rates for grass weed control. If Nuquat 250 has been applied use a suitable post-emergent grass herbicide (such as Nufarm Asset) at recommended label rates after regrowth but before heading.
		NSW only	1.2L/ha	
	Annual grass and some broadleaf weeds.	Qld, SA, Tas, Vic, NT, WA only	1.6-2.4L/ha <sup>‡</sup>	
		NSW, ACT only	1.2L/ha	
Perennial Grass Seed Crops  Cocksfoot, perennial ryegrass Phalaris and Demeter Fescue only	Annual grass and some broadleaf weeds	All States	600mL-1.2L/ha <sup>‡</sup>	Use the low rate for cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter Fescue.  Spray after about 4 weeks after a full weed germination following the Autumn break. The perennial grasses must be at least 12 months old at spraying.
Spray topping to reduce seed set Chickpeas Faba beans Field peas Lentils Lupins Vetch	Annual ryegrass	All States	400 or 800mL/ha	<b>As an aid in managing annual ryegrass resistance. For use on escapes from a previous herbicide application in the current crop.</b> Spray the crop when the ryegrass is at the optimum stage, that is when the last ryegrass seed heads at the bottom of the plant have emerged and the majority are at or just past flowering (with anthers present or glumes open) but before haying off is evident - usually October to November. Use of the higher rate in these crops is usually more reliable and gives a greater reduction in seed set. Reduction in crop yield may occur especially if the crop is less advanced relative to the ryegrass that is if crops have a majority of green immature pods. The higher rate may also increase any yield reduction. In practice crop losses in excess of 25% may occur. Apply by ground boom only in 50-100L/ha. Spray with a calibrated boom spray raised to give double overlap at the level of the ryegrass seed heads. Pressures of 250-350kPa and use of 110015 or 02 nozzles or equivalent will aid coverage.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Spray topping to reduce seed set  Pastures	Grasses generally (particularly annual ryegrass)	All States	400mL/ha	Heavily graze paddocks during Spring flush to encourage even head development. Remove stock 2-3 weeks before the anticipated maturity date of the target species. However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed heads at the bottom of the plant have emerged and initial signs of haying off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Barley grass			Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening of grass seeds, stock (excepting horses) may be returned 24 hours after spraying. Where hardening seeds are present harrow to knock seed from the heads. DO NOT introduce lambs into paddock until safe from risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with calibrated boom spray raised to give double overlap at the level of the seed heads.
	Saffron thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.
Prevention of annual ryegrass toxicity	Spray top - Graze to destroy seed heads	WA only	400mL/ha	Grazing management as for spray topping above. Remove stock 3-4 weeks before the anticipated maturity date. Spray must be applied within 10 days after emergence of the first ryegrass seed heads. To ensure adequate control of toxin development, <b>heavy continuous grazing is essential from 1 day after spraying</b> until the pasture has completely hayed off. The required stocking rate will vary but must be sufficient to keep all regrowth after spraying completely eaten off to prevent further growth producing new seed heads which could become toxic
Hay freezing	Maximum retention of protein in standing dry feed	All States	800mL/ha	Graze paddocks as for spray topping above. Remove stock 3-4 weeks before the anticipated maturity date. Apply prior to commencement of haying off regardless of the grass species involved. Spray with calibrated boom spray raised to give double overlap at the level of the seed heads.
General Weed Control Bananas	Annual weed control	Qld, NSW, WA, NT only	1.6-3.2L sprayed ha <sup>Ψ</sup>  160-320mL/100L  Misters 8mL/L	Spray as necessary for control of annual weeds. Avoid contacting crop foliage. Apply soon after weed emergence and before weeds reach 15cm in height. Use spraying pressure less than 240kPa. Avoid chemical contact with roots and peepers near the pseudo stem. Repeat sprays as required.
Hops	Annual grasses	Vic, Tas only	1.2-1.6L/ha plus 1.1kg/ha Nufarm Simazine 900 DF and/or 750mL-1.4L/ha Reglone <sup>Ψ</sup>	Apply as a directed inter-row spray prior to crop emergence from Winter dormancy, using a minimum of 250L/ha spray volume to ensure good and even coverage of weeds.
Orchards Vineyards	Annual weed control	Qld, Vic, SA, WA, Tas, NT only	1.6-3.2L/sprayed ha <sup>Ψ</sup>	Spray as necessary for control of annual weeds. Avoid contacting crop foliage. Nuquat 250 will not harm trees or vines with mature brown bark if this alone is sprayed. Use the higher rate for dense weed growth.
		NSW only	1.7L/sprayed ha <sup>Ψ</sup>	If fat hen <i>Chenopodium album</i> or <i>Portulaca</i> spp. are present and Nuquat 250 rate is less than the ratio 800mL/100L add 140mL Activator or 120mL Chemwet 1000 per 100L of spray mix.
Peanuts Post-emergence (in crop)	Datura spp. (2 to 4 leaf)	Qld, NT only	400mL /ha	Spray peanuts up to 7-8 leaf stage but before majority of plants are flowering. Foliage will be scorched following application but plants recover rapidly. Apply in 200-250L/ha for thorough coverage of weed foliage. A dense canopy of weeds may reduce weed control due to shielding. Add 70mL Activator or 60mL Chemwet 1000 per 100L of spray mix.  DO NOT spray (on peanuts) under extremely hot dry conditions when peanuts are very small. In environments such as Far North Queensland use the lower rates in the range.
	Annual ground cherry (2 to 3 leaf)		600mL/ha	
	Apple-of-Peru (2 to 4 leaf)		600mL/ha	
	Milkweed (2 to 3 leaf)		600mL/ha	
	Stagger weed (2 to 3 leaf)		800mL/ha	
	Blue heliotrope (2 to 3 leaf)		800mL/ha	
	Wandering Jew (2 to 3 leaf)		800mL/ha	
	Anoda weed (2 to 3 leaf)		800mL/ha	
	Bellvine (2 to 3 leaf)		1L/ha	
Common morning glory (2 leaf)	1L/ha			

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Potatoes	General weed control (in-crop)	All States	1.2-1.6L/ha <sup>ψ</sup>	Spray at early crop emergence (no later than 25% emergence of potato shoots). Use the higher rate for dense weed growth.
	Pre-harvest weed control		2.8L/ha <sup>ψ</sup>	Spray about one week before digging and after tops have died down.
Row Crops, Vegetables and Market Gardens	Pre-planting and pre-crop emergence	All States	1.2-1.6L/ha OR 200mL/100L <sup>ψ</sup> <sup>^</sup>	To control weeds in seed beds. Treat no less than three days before sowing or before crop emergence. Use the lower rate for early Autumn applications.
	Post-emergence inter-row weed control		1.2-1.6L/ha OR 200mL/100L <sup>ψ</sup> <sup>^</sup>	Apply after crop seedlings have emerged or when transplanted crops are established. Direct the spray so that it does not touch the crop. Use shielded nozzles
	Seedling weeds			Seedling weeds - use the lower rate for early Autumn applications.
	Older weeds		2.4L or 400mL/100L <sup>ψ</sup>	More mature stages of weed growth.
Non-Agricultural situations, around sheds, roadways, paths	Annual weed control	All States	1.6- 4L/ha OR 200mL/100L <sup>ψ</sup> <sup>^</sup>	Spray to thoroughly wet weed growth. Nuquat 250 can be combined with soil residual herbicides Nufarm Diuron 900DF or Nufarm Simazine 900DF to give rapid knockdown and prolonged weed control. Use the higher rate for dense weed growth
	Columbus grass	NSW only	<sup>^</sup> <b>Spot Spraying</b> 160mL/100L plus 1L of a 745g/L flupropanate product <b>Boomspray</b> 2.3-4.5L/ha plus 12-22L of a 745g/L flupropanate product	
Firebreaks	Knock down weed growth to eliminate fire hazard or assist firebreak burn	All States	1.6L-4L/ha	Apply mid-Winter to early Summer. Use the higher rate for dense weed growth. After desiccation is complete the sprayed area may be burnt (normally 7-10 days after spraying). Nuquat 250 can be combined with soil residual herbicides Nufarm Diuron 900DF or Flowable Diuron Herbicides or Nufarm Simazine 900DF or Nufarm Nu-Trazine 900 DF or Flowable Nu-Trazine Herbicides to give rapid knockdown and prolonged weed control.

CROP	WEEDS	GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Sugar Cane (Plant and ratoon)	Grass and some broadleaf weeds	up to 5cm high	Qld, NSW, NT only	1.2-1.6L/ sprayed ha	Apply as a broadcast spray over-the-top of plant cane up to the 3-4 leaf stage or ratoon cane up to 10cm high. Cane foliage will be scorched but new leaves will appear in 7-10 days. In plant cane between the 3-4 leaf stage and the formation of the true stem use a directed interspace spray. The Irvin spray boom (or other similar equipment) is the most suitable equipment to avoid excessive drift onto cane foliage while spraying at the bases of plant and ratoon cane. After the formation of the true stem which is resistant to Nuquat 250, the sprayer height can be raised to overlap the spray pattern to give weed control in the stool. Use the higher rate for dense, more mature weeds. Nuquat 250 can be mixed with Nufarm Nu-Trazine 900 DF or Nufarm Flowable Nu-Trazine herbicides to give residual weed control when used as a blanket or directed spray - refer to the Nufarm Nu-Trazine 900 DF or Flowable Nu-Trazine label for specific rates. It may also be mixed with Nufarm Diuron 900DF at 2.8-3.9kg/ha or Nufarm Flowable Diuron at 5L or 7L/ha for residual weed control <sup>^</sup> To enhance activity of Nuquat 250 under favourable growing conditions and in open sunny conditions add Nufarm Diuron 900DF or Flowable Diuron at rates shown for weed size. Nufarm Diuron 900DF at rates up to 500g/ha or Flowable Diuron at rates up to 900mL/ha can be blanket sprayed. Use a directed spray for higher rates of Nufarm Diuron 900DF or Flowable Diuron. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5cm high use a minimum of 250L spray solution/ha, increase to 350L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds >10cm high and for dense mature weeds. Always add Activator at 140mL or Chemwet 1000 at 120mL per 100L of water.
	Grass and some broadleaf weeds - enhancement with Nufarm Diuron 900DF or Flowable Diuron	up to 5cm high		1.2-1.6L/ha + 275-500g Nufarm Diuron 900DF or 500-900mL Flowable Diuron	
	Grass and some broadleaf weeds - enhancement with Nufarm Diuron 900DF or Flowable Diuron	up to 10cm high		1.2-1.6L/ha + 1kg Nufarm Diuron 900DF or 1.8L Flowable Diuron	
	Grass and some broadleaf weeds - enhancement with Nufarm Diuron 900DF or Flowable Diuron and short residual control	>10cm high		1.6L/ha + 2.8 to 3.9kg Nufarm Diuron 900DF or 5 to 7L Flowable Diuron	

<sup>ψ</sup> Capeweed or Erodium spp. present: Add Reglone at 750mL-1.5L/ha (125mL or 250mL/100L for high volume spraying). Use higher rate for plants more than 10cm diameter.

<sup>^</sup> If Nuquat 250 rate is less than 400mL/100L add 60mL Activator or 100mL Chemwet 1000 per 100L of spray mix.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE. THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.**

**WITHHOLDING PERIOD:**

**DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR AT LEAST 1 DAY, OR GRAZE HORSES FOR 7 DAYS AFTER APPLICATION.**

**REMOVE STOCK FROM TREATED AREAS 3 DAYS BEFORE SLAUGHTER.**

**CHICK PEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS AND VETCH - DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.**

**PLEASE NOTE EXTRA WETTER REQUIREMENTS FOR HIGH VOLUME SPRAYING.**

**RESISTANT WEEDS WARNING**

GROUP	L	HERBICIDE
-------	---	-----------

Nufarm Nuquat 250 Non-Residual, Knockdown Herbicide ("Nuquat 250") is a member of the Bipyridyls group of herbicides. Nuquat 250 has the inhibitors of photosynthesis at photosystem I mode of action. For weed resistance management Nuquat 250 is a Group L Herbicide. Some naturally-occurring weed biotypes resistant to Nuquat 250 and other Group L herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nuquat 250 or other Group L herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nuquat 250 to control resistant weeds.

This product kills annual grasses and most annual broadleaf weeds (excluding capeweed) in specified situations and should not be used for any other purpose. Quickly kills green plant tissue on contact. Nuquat 250 is immediately inactivated in the soil. At spraying, weeds should be growing vigorously and must not be covered with soil or heavy dew. The principle of selective weed control with this product is that annual weeds are killed but perennial plants and clovers recover after an initial scorch. The control of annual weeds by spraying with this product will allow the desirable perennial species to thicken up at the expense of the weeds. Moisture and fertility should not be limiting at spraying and the proportion of desirable species must be great enough for them to fill in the areas previously occupied by weeds. Long term weed control can be obtained following the quick knockdown given by this product if it is combined with soil residual chemicals.

1. DO NOT use hand-held ultra low volume controlled droplet applicators (CDA units), boomless jets or misting-machines (except in banana plantations).

**2. MIXING**

Add the required quantity of product to water in the spray tank and agitate to give even mixing. Agitate again if left standing.

**3. WETTING AGENT**

This product contains a wetting agent and additional wetter is not required unless high volume spraying results in excessive dilution of wetter content. This will occur when product rates fall below 400mL per 100L of spray. Under such circumstances wetter should be added at the rate of 70mL of Activator or 100mL of Chemwet 1000 per 100L of spray mix. Where Fat Hen or Portulaca are present in orchard or vineyard situations, extra wetter should be used when this product ratio is less than 800mL per 100L. Add wetter at double the above recommendations. DO NOT use alkaline or anionic wetting agents.

**4. CLEAN WATER**

Mix this product **with clean water only**. Water should be clean and free from clay, silt and algae. Providing it meets this requirement, saline water, water collected from roofs, bore water, dam water and water from creeks may be used.

**5. APPLICATION****(i) Cereals and Broadacre Spraying**

Use only through a properly calibrated boom spray which should be fitted with flat fan jets and adjusted to a height to give at least double overlap of the spray at the top of the weeds being sprayed. It is essential that a good marking system be used. If a disc marker is used, it must be mounted so as to turn the soil back on to the area sprayed. It is essential to obtain good leaf coverage with the spray and volumes of dilute spray must be adjusted according to density of weed growth. 100L/ha may be used for seedlings or well grazed weeds up to 2cm high. For plant height 2-5 cm use 150L/ha and up to 6-10cm use 200L/ha. Spray volumes may be as low as 50L/ha (30L/ha in WA) for weed growth below 5cm high, or for spray topping and hay freezing. Equipment must be appropriate to this volume, properly calibrated and fitted with spraying tips designed to give a medium spray quality (ASAE S572) at the target.

**(ii) High Volume Application**

Higher volumes will generally be required to give good coverage of weed growth in situations other than those specified under cereals and other broadacre crops.

**(iii)** Wash spray equipment with clean water immediately after use. This product is highly corrosive to metals, particularly galvanised iron and aluminium and should not be left for long periods in tanks or equipment made of these materials.

**FOR GROUND APPLICATION ONLY**

DO NOT use this formulation through aircraft, misting machines (except in banana plantations) or hand-held ultra low volume controlled droplet applicators (CDA units).

**6. COMPATIBILITY**

This product combines satisfactorily with the soil active herbicides Nufarm Nu-trazine 900DF and Flowable Nu-Trazine, Nufarm Diuron 900DF and Flowable Diuron and Nufarm Simazine 900DF and Flowable Simazine where prolonged weed control is required as well as a quick knockdown. This product is compatible with Activator®, Chemwet 1000, Reglone\*, Revolver®, Spray.Seed\* 250, Kamba® 500, Amicide® 625, Nufarm Surpass 475, Kamba® M, MCPA 500 (no more than 1L/800mL Nuquat® 250), Glean\*, Lusta®, Yield\*, Avadex® Xtra, TriflurX® and Striker® (oxyfluorfen).

**7. SPRAYING CONDITIONS**

Avoid spraying plants under stress from waterlogging, frost, drought, etc. or covered with dust and soil. Results will be better if application is made in dull weather or at the end of the day. Light rain following spraying will not affect results. Avoid drift into neighbouring crops.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. This formulation should not be applied on or near water which is used for irrigation purposes.

**PROTECTION OF LIVESTOCK**

Domestic pets and poultry - keep away from treated areas. This formulation should not be applied on or near water which is used for livestock watering.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. This formulation should not be applied on or near water which is used for human consumption, livestock watering or irrigation purposes or water used for commercial or recreational fishing.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should NOT be burnt.

**Envirodrum 110L Mini Bulk Returnable Container**

Store in the original sealed Envirodrum in a cool, well ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

**(1000L)**

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks eyes. Protect eyes while using. Avoid contact with eyes, skin and clothing. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use wash contaminated clothing, gloves and face shield or goggles.

**SPRAY APPLICATION**

- DO NOT work in spray mist.
- DO NOT continue to use if skin irritation or nose bleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
- When using misting machines for weed control in banana plantations, cut back to run at half throttle, thus preventing the production of fine droplets, the inhalation of which may be dangerous.
- When there is a risk of exposure to spray mist or when using misting machines in banana plantations wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended, but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
- Avoid contacting vegetation wet with spray, but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

**FIRST AID**

If poisoning occurs, get to a doctor or hospital quickly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**NOTE TO PHYSICIANS.**

For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning; A Guide For Doctors".

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited ACN 004 377 780  
103-105 Pipe Road Laverton North Victoria 3026  
Tel: (03) 9282 1000 Fax: (03) 9282 1001

© Nuquat, Kamba, Amicide, TriflurX, Surpass, Lusta, Revolver and Asset are Registered Trade Marks of Nufarm Australia Limited.

© Avadex and Striker are registered trademarks of Nufarm Technologies USA Pty Ltd

© Activator is a registered trademark of Loveland Industries Inc.

\*Other registered trademarks